


IN THE CLAIMS

Please amend the claims as follows:


1. (Currently Amended) A compound of formula



wherein

R is a hydrogen atom or a methyl group;

R₁ is a hydrogen atom, an N,N-dimethylamino-N-oxide group, an N-benzyl-N-methylamino group, an N-acetyl-N-methylamino group, an N-[N,N-dimethylamino(C₁-C₂)alkylamino]acetyl-N-methylamino group or a chain of formula



wherein

A is a hydrogen atom, a phenyl or a five- or six-membered heteroaryl ring selected from pyrrole, thiophene, furan, imidazole, oxazole, thiazole, pyridine, pyrimidine, triazole and thiadiazole;

X is O or NR₆ and R₆ is a hydrogen atom;

Y is, when n is 0, a C₆H₄ group or a five- or six-membered heteroaryl ring selected from pyrrole, thiophene, furan, imidazole, oxazole, thiazole, pyridine, pyrimidine, triazole and thiadiazole; or, when n is 1, NR₆ and R₆ is a hydrogen atom;

r is an integer from 1 to 3;


m is the integer 1 or 2;

n is the integer 0 or 1;

or R₁ forms a bond together with R₂

~~R₁ is a hydrogen atom, an N,N-di(C₁-C₃)alkylamino group, an N,N-di(C₁-C₃)alkylamino-N-oxide group, an N-(C₁-C₃)alkyl-N-benzyl-amino group, an N-(C₁-C₄)acyl-N-(C₁-C₃)alkylamino group, an N-[N,N-dimethylamino(C₁-C₄)alkylamino]acetyl-N-(C₁-C₃)alkylamino group~~

~~or a chain of formula~~



~~wherein~~

~~A is a hydrogen atom, a phenyl or a five- or six- membered heteroaryl ring having from one to three hetero atoms selected from nitrogen, oxygen and sulphur;~~

~~X is O, S, SO, SO₂ or NR₆, where R₆ is a hydrogen atom, a linear or branched C₁-C₃ alkyl, a C₁-C₃ alkoxy carbonyl group or a benzyloxy carbonyl group;~~

~~Y is a C₆H₄ group, a five- or six- membered heteroaryl ring having from one to three hetero atoms selected from nitrogen, oxygen and sulphur or is O, S, SO, SO₂ or NR₆ where R₆ has the meanings given above;~~

~~r is an integer from 1 to 3;~~


~~m is an integer from 1 to 6;~~

~~n is an integer from 0 to 2;~~

~~or R₁ forms a bond together with R₂;~~

~~R₂ is a hydrogen atom or forms a bond together with R₁;~~

~~R₃ is a hydroxy group or forms a group =N-O-R₅ together with R₄, and R₅ is a hydrogen atom, a linear or branched C₁-C₅ alkyl, a benzyl optionally substituted with one or two substituents selected from nitro, hydroxy, carboxy, amino, linear or branched C₁-C₅ alkyl, C₁-C₄ alkoxy carbonyl groups, aminocarbonyl groups or cyano groups or a chain of formula~~



~~wherein~~

~~r, m, n, X, Y and A have the meanings given above;~~

R_4 is a hydrogen atom or forms a group $=N-O-R_5$ together with R_3 , and R_5 has the meanings given above;

and the pharmaceutically acceptable salts thereof;


~~provided, however, that~~

~~R_1 is not a dimethylamino group when R_3 is hydroxy, and both R_2 and R_4 are a hydrogen atom;~~

~~R_1 is not a dimethylamino group when in the substituent $=N-O-R_5$ in the 9 position, R_5 is a hydrogen atom, a linear or branched C_1-C_5 -alkyl, an unsubstituted benzyl group, or a chain $-(CH_2)_r-X-(CH_2)_m-Y-(CH_2)_n-A$ where r is 1, X is O , m is 2, Y is O , n is 1, and A is H ;~~

~~R_1 is not a methylethylamino group when in the substituent $=N-O-R_5$ in the 9 position, R_5 is a linear or branched C_1-C_5 -alkyl, or an unsubstituted benzyl group.~~

2. (Original) A compound according to Claim 1, wherein the oxime group that may be present in position 9 is of E configuration.
3. (Cancelled)
4. (Cancelled)
5. (Currently Amended) A compound according to Claim 1, ~~Claim 4~~, wherein R_1 is a hydrogen atom, an N,N-dimethylamino-N-oxide group, an N-benzyl-N-methylamino group, an N-acetyl-N-methylamino group, an N-[N,N-dimethylaminoethylamino]acetyl-N-methylamino group or a chain of formula



wherein

A is a hydrogen atom, a phenyl or a heteroaryl ring selected from thiophene, furan, thiazole, pyridine and triazole;


X is NR_6 and R_6 is a hydrogen atom;

Y is, when n is 0, a C₆H₄ group or a heteroaryl ring selected from thiophene, furan, thiazole, pyridine and triazole; or, when n is 1, NR₆ and R₆ is a hydrogen atom; or R₁ forms a bond together with R₂.

6. (Cancelled)

7. (Cancelled)


8. (Currently Amended) ~~A compound according to Claim 7,~~ A compound of formula (I):



wherein

R is a hydrogen atom or a methyl group;

R₁ is a hydrogen atom, an N,N-dimethylamino-N-oxide group, an N-benzyl-N-methylamino group, an N-acetyl-N-methylamino group, an N-[N,N-dimethylamino(C₁-C₂)alkylamino]acetyl-N-methylamino group or a chain of formula



wherein

A is a hydrogen atom, a phenyl or a five- or six-membered heteroaryl ring selected from pyrrole, thiophene, furan, imidazole, oxazole, thiazole, pyridine, pyrimidine, triazole and thiadiazole;

X is O or NR₆ and R₆ is a hydrogen atom;

Y is, when n is 0, a C₆H₄ group or a five- or six-membered heteroaryl ring selected from pyrrole, thiophene, furan, imidazole, oxazole, thiazole, pyridine, pyrimidine, triazole and thiadiazole; or, when n is 1, NR₆ and R₆ is a hydrogen atom;

r is an integer from 1 to 3;

m is the integer 1 or 2;

n is the integer 0 or 1;

or R₁ forms a bond together with R₂;


R₂ is a hydrogen atom or forms a bond together with R₁;

R₃ is a hydroxy group;

R₄ is a hydrogen atom;

and the pharmaceutically acceptable salts thereof.

9. (Original) A compound according to Claim 8, wherein R₁ is a hydrogen atom, an N,N-dimethylamino-N-oxide group, an N-benzyl-N-methylamino group, an N-acetyl-N-methylamino group, an N-[N,N-dimethylaminoethylamino]acetyl-N-methylamino group or a chain of formula



wherein

A is a hydrogen atom, a phenyl or a heteroaryl ring selected from thiophene, furan, thiazole, pyridine and triazole;


X is NR₆ and R₆ is a hydrogen atom;

Y is, when n is 0, a C₆H₄ group or a heteroaryl ring selected from thiophene, furan, thiazole, pyridine and triazole; or, when n is 1, NR₆ and R₆ is a hydrogen atom;

or R₁ forms a bond together with R₂.


10. - 16. (Cancelled)

17. (Original) A process for preparing a compound according to Claim 1, characterized in that the L-cladinose moiety in 3 position is removed from the erythromycin A compounds of formula



wherein R, R₁, R₂, R₃ and R₄ are defined as in Claim 1;
via a hydrolysis reaction.

18. (Original) Process according to Claim 17, wherein in formula II R_3 is a hydroxyl group and R_4 is a hydrogen atom.
19. (Original) Process according to Claim 17, wherein the removal of the cladinose is performed via an acid hydrolysis reaction catalyzed in the presence of a mineral acid and a protic organic solvent.
20. (Currently Amended) A compound of formula II:




wherein

R is a hydrogen atom or a methyl group;

R₁ is a hydrogen atom, an N,N-di(C₁-C₃)alkylamino group, an N,N-di(C₁-C₃)alkylamino-N-oxide group, an N-(C₁-C₃)alkyl-N-benzylamino group, an N-(C₁-C₄)acyl-N-(C₁-C₃)alkylamino group, an N-[N,N-dimethylamino(C₁-C₄)alkylamino]acetyl-N-(C₁-C₃)alkylamino group

or a chain of formula



wherein

A is a hydrogen atom, a phenyl or a five- or six-membered heteroaryl ring having from one to three hetero atoms selected from nitrogen, oxygen and sulphur;

X is O, S, SO, SO₂ or NR₆, where R₆ is a hydrogen atom, a linear or branched C₁-C₃ alkyl, a C₁-C₃ alkoxy carbonyl group or a benzyloxy carbonyl group;

Y is a C₆H₄ group, a five- or six-membered heteroaryl ring having from one to three hetero atoms selected from nitrogen, oxygen and sulphur or is O, S, SO, SO₂ or NR₆ where R₆ has the meanings given above;

r is an integer from 1 to 3;

m is an integer from 1 to 6;

n is an integer from 0 to 2;

or R₁ forms a bond together with R₂;

R₂ is a hydrogen atom or forms a bond together with R₁;
R₃ is a hydroxy group;
R₄ is a hydrogen atom;
and the pharmaceutically acceptable salts thereof;
provided, however, that (i) R₁ is not an N,N-dimethyl amino group, and (ii) R₁ is not an N,N-dimethyl amino-N-oxide group when R is a hydrogen atom,
and provided that one of the following limitations is met:

R is a hydrogen atom and R₁ forms a bond together with R₂;

R is a hydrogen atom and R₁ is an N-benzyl-N-methylamino group;

R is a hydrogen atom and R₁ is an N-acetyl-N-methylamino group;

R is a hydrogen atom and R₁ is an N-[N,N-dimethylaminoethylamino]acetyl-N-methyl amino group;


R is a hydrogen atom and R₁ is an N-methyl-N-3-[(2-thiazolylmethyl)amino]propylamino group;

R is a hydrogen atom and R₁ is an N-2-[2-[(2-thiazolylmethyl)amino]ethylamino]ethyl-N-methylamino group; and

R is a hydrogen atom and R₁ is an N-2-[2-(benzylamino)ethylamino]ethyl-N-methylamino group.

21. (Original) A compound according to Claim 20, wherein R is a hydrogen atom and R₁ forms a bond together with R₂.
22. (Original) A compound according to Claim 20, wherein R is a hydrogen atom and R₁ is an N-benzyl-N-methylamino group.
23. (Original) A compound according to Claim 20, wherein R is a hydrogen atom and R₁ is an N-acetyl-N-methylamino group.
24. (Original) A compound according to Claim 20, wherein R is a hydrogen atom and R₁ is an N-[N,N-dimethylaminoethylamino]acetyl-N-methyl amino group.
25. (Original) A compound according to Claim 20, wherein R is a hydrogen atom and R₁ is an N-methyl-N-3-[(2-thiazolylmethyl)amino]propylamino group.

26. (Original) A compound according to Claim 20, wherein R is a hydrogen atom and R₁ is an N-2-[2-[(2-thiazolylmethyl)amino]ethylamino]ethyl-N-methylamino group.
27. (Original) A compound according to Claim 20, wherein R is a hydrogen atom and R₁ is an N-2-[2-(benzylamino)ethylamino]ethyl-N-methylamino group.
28. (Previously Presented) The compound de(N-methyl)-9-dihydroerythromycin A.
29. (Previously Presented) The compound de(N-methyl)-descladinosyl-9-dihydroerythromycin A.
30. (Cancelled)
31. (Cancelled)
32. (Cancelled)
33. (New) A process for preparing a compound according to Claim 8, wherein the L-cladinosyl moiety in 3 position is removed from the erythromycin A compounds of formula



wherein R, R₁, R₂, R₃ and R₄ are defined as in Claim 8;
via a hydrolysis reaction.

34. (Original) Process according to Claim 33, wherein the removal of the cladinose is performed via an acid hydrolysis reaction catalyzed in the presence of a mineral acid and a protic organic solvent.